-1-

A method for inhibiting cyclooxygenase or prostaglandin H synthase enzymes which comprises:

providing at least one compound isolatable from a cherry with at least one of the enzymes to inhibit the enzymes.

-2-

A method for inhibiting cyclooxygenase or prostaglandin H synthase enzymes which comprises:

providing at least one bioflavonoid compound isolatable from a cherry with at least one of the enzymes to inhibit the enzymes.

-3-

The method of Claim 1 wherein the method is in vitro.

-4-

The method of Claim 1 wherein the method is  $in\ vivo.$ 

-5-

The method of any one of Claims 1, 2, 3 or 4 wherein the compound is from a tart cherry.

-6-

The method of any one of Claims 1, 2, 3 or 4 wherein the compound is from a sweet cherry.

The transfer of the transfer o

5

10

15

20

25

A method for inhibiting inflammation in a mammal which comprises:

administering at least one compound isolated from a cherry to the mammal to inhibit inflammation.

-8-

The method of Claim 7 wherein the mammal is human.

10

5

-9-

The method of any one of Claims 7 or 8 wherein the compound is from a tart cherry.

15

-10-

The method of Claim 7 wherein the compound is from a sweet cherry.

-11-

20

A method for inhibiting inflammation in a mammal which comprises:

administering at least one bioflavonoid compound isolatable from a cherry to the mammal to inhibit the inflammation.

25

-12-

 $\,$  The method of Claim 11 wherein the compound is from a tart cherry.

30

-13-

The method of Claim 11 wherein the compound is from a sweet cherry.

-14-

35

The method of Claim 11 wherein the mammal is human.

The method of Claim 1 wherein the compound is contained in a composition which comprises a dried mixture of isolated anthocyanins, bioflavonoids and phenolics from cherries and a food grade carrier.

-16-

The method of Claim 15 wherein the carrier is dried cherry pulp.

10

5

-17-

The method of Claim 15 wherein the ratio of dried mixture to carrier is between about 0.1 to 100 and 100 to 0.1.

15

-18-

The method of any one of Claims 1, 2, 3 or 4 wherein the compound is incorporated into a food.

20

25

3.0

-19-

The method of Claim 7 wherein the compound is incorporated into a food.

-20-

The method of Claim 11 wherein the bioflavonoid is incorporated into a food.

-21-

A method for inhibiting inflammation in a mammal which comprises:

administering anthocyanin including cyanidin to the mammal to inhibit inflammation.

-22-

35 The method of Claim 21 wherein the anthocyanin is isolated from a tart cherry.

The state of the s

The method of Claim 21 wherein the anthocyanin is isolated from a sweet cherry.

-24-

The method of Claim 21 wherein the mammal is human.

-25-

The method of any one of Claims 21, 22, 23 or 24 wherein the anthocyanin is incorporated into a food.

-26-

The method of Claim 21 wherein the anthocyanin is essentially pure.

The property of the control of the c

5

15